

CHAIN ENUMERATION OF k -DIVISIBLE NONCROSSING PARTITIONS OF CLASSICAL TYPES

JANG SOO KIM

ABSTRACT. We give combinatorial proofs of the formulas for the number of multichains in the k -divisible noncrossing partitions of classical types with certain conditions on the rank and the block size due to Krattenthaler and Müller. We also prove Armstrong's conjecture on the zeta polynomial of the poset of k -divisible noncrossing partitions of type A invariant under the 180° rotation in the cyclic representation.

1. INTRODUCTION

For a finite set X , a *partition* of X is a collection of mutually disjoint nonempty subsets, called *blocks*, of X whose union is X . Let $\Pi(n)$ denote the poset of partitions of $[n] = \{1, 2, \dots, n\}$ ordered by refinement, i.e. $\pi \leq \sigma$ if each block of σ is a union of blocks of π . There is a natural way to identify $\pi \in \Pi(n)$ with an intersection of reflecting hyperplanes of the Coxeter group A_{n-1} . For this reason, we will call $\pi \in \Pi(n)$ a *partition of type A_{n-1}* . With this observation Reiner [11] defined partitions of type B_n and type D_n as follows. A *partition of type B_n* is a partition π of $[\pm n] = \{1, 2, \dots, n, -1, -2, \dots, -n\}$ such that if B is a block of π then $-B = \{-x : x \in B\}$ is also a block of π , and there is at most one block, called *zero block*, which satisfies $B = -B$. A *partition of type D_n* is a partition of type B_n such that its zero block, if exists, has more than two elements. Let $\Pi_B(n)$ (resp. $\Pi_D(n)$) denote the poset of type B_n (resp. type D_n) partitions ordered by refinement.

A *noncrossing partition of type A_{n-1}* , or simply a noncrossing partition, is a partition $\pi \in \Pi(n)$ with the following property: if integers a, b, c and d with $a < b < c < d$ satisfy $a, c \in B$ and $b, d \in B'$ for some blocks B and B' of π , then $B = B'$.

Let k be a positive integer. A noncrossing partition is called *k -divisible* if the size of each block is divisible by k . Let $\text{NC}(n)$ (resp. $\text{NC}^{(k)}(n)$) denote the subposet of $\Pi(n)$ (resp. $\Pi(kn)$) consisting of the noncrossing partitions (resp. k -divisible noncrossing partitions).

Bessis [4], Brady and Watt [5] defined the generalized noncrossing partition poset $\text{NC}(W)$ for each finite Coxeter group W , which satisfies $\text{NC}(A_{n-1}) \cong \text{NC}(n)$. Armstrong [1] defined the poset $\text{NC}^{(k)}(W)$ of generalized k -divisible noncrossing partitions for each finite Coxeter group W , which reduces to $\text{NC}(W)$ for $k = 1$ and satisfies $\text{NC}^{(k)}(A_{n-1}) \cong \text{NC}^{(k)}(n)$.

Date: October 11, 2009.

2000 Mathematics Subject Classification. 05A15; 05E15.

Key words and phrases. noncrossing partitions, chain enumeration.

For each classical Coxeter group W , we have a combinatorial realization of $\text{NC}^{(k)}(W)$. In other words, similar to $\text{NC}(n)$ and $\text{NC}^{(k)}(n)$, there are combinatorial posets $\text{NC}_B(n) \subset \Pi_B(n)$, $\text{NC}_B^{(k)}(n) \subset \Pi_B(kn)$, $\text{NC}_D(n) \subset \Pi_D(n)$ and $\text{NC}_D^{(k)}(n) \subset \Pi_D(kn)$, which are isomorphic to $\text{NC}(B_n)$, $\text{NC}^{(k)}(B_n)$, $\text{NC}(D_n)$ and $\text{NC}^{(k)}(D_n)$ respectively. Reiner [11] defined the poset $\text{NC}_B(n)$ of noncrossing partitions of type B_n , which turned out to be isomorphic to $\text{NC}(B_n)$. This poset is naturally generalized to the poset $\text{NC}_B^{(k)}(n)$ of k -divisible noncrossing partitions of type B_n . Armstrong [1] showed that $\text{NC}_B^{(k)}(n) \cong \text{NC}^{(k)}(B_n)$. Athanasiadis and Reiner [3] defined the poset $\text{NC}_D(n)$ of noncrossing partitions of type D_n and showed that $\text{NC}_D(n) \cong \text{NC}(D_n)$. Krattenthaler [9] defined the poset $\text{NC}_D^{(k)}(n)$ of the k -divisible noncrossing partitions of type D_n using annulus and showed that $\text{NC}_D^{(k)}(n) \cong \text{NC}^{(k)}(D_n)$; see also [8].

In this paper we are mainly interested in the number of multichains in $\text{NC}^{(k)}(n)$, $\text{NC}_B^{(k)}(n)$ and $\text{NC}_D^{(k)}(n)$ with some conditions on the rank and the block size.

Definition 1. For a multichain $\pi_1 \leq \pi_2 \leq \cdots \leq \pi_\ell$ in a graded poset P with the maximum element $\hat{1}$, the *rank jump vector* of this multichain is the vector $(s_1, s_2, \dots, s_{\ell+1})$, where $s_1 = \text{rank}(\pi_1)$, $s_{\ell+1} = \text{rank}(\hat{1}) - \text{rank}(\pi_\ell)$ and $s_i = \text{rank}(\pi_i) - \text{rank}(\pi_{i-1})$ for $2 \leq i \leq \ell$.

We note that all the posets considered in this paper are graded with the maximum element, however, they do not necessarily have the minimum element. We also note that the results in this introduction have certain ‘obvious’ conditions on the rank jump vector or the block size, which we will omit for simplicity.

Edelman [6, Theorem 4.2] showed that the number of multichains in $\text{NC}^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ is equal to

$$(1) \quad \frac{1}{n} \binom{n}{s_1} \binom{kn}{s_2} \cdots \binom{kn}{s_{\ell+1}}.$$

Modifying Edelman’s idea of the proof of (1), Reiner found an analogous formula for the number of multichains in $\text{NC}_B(n)$ with given rank jump vector. Later, Armstrong generalized Reiner’s idea to find the following formula [1, Theorem 4.5.7] for the number of multichains in $\text{NC}_B^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$:

$$(2) \quad \binom{n}{s_1} \binom{kn}{s_2} \cdots \binom{kn}{s_{\ell+1}}.$$

Athanasiadis and Reiner [3, Theorem 1.2] proved that the number of multichains in $\text{NC}_D(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ is equal to

$$(3) \quad 2 \binom{n-1}{s_1} \binom{n-1}{s_2} \cdots \binom{n-1}{s_{\ell+1}} + \sum_{i=1}^{\ell+1} \binom{n-1}{s_1} \cdots \binom{n-2}{s_i-2} \cdots \binom{n-1}{s_{\ell+1}}.$$

To prove (3), they [3, Lemma 4.4] showed the following using incidence algebras and the Lagrange inversion formula: the number of multichains $\pi_1 \leq \pi_2 \leq \cdots \leq \pi_\ell$ in $\text{NC}_B(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ such that i is the smallest integer for which π_i has a zero block is equal to

$$(4) \quad \frac{s_i}{n} \binom{n}{s_1} \binom{n}{s_2} \cdots \binom{n}{s_{\ell+1}}.$$

Since (4) is quite simple and elegant, it deserves a combinatorial proof. In this paper we prove a generalization of (4) combinatorially; see Lemma 15.

The number of noncrossing partitions with given block sizes has been studied as well. In the literature, for instance [1, 2, 3], $\text{type}(\pi)$ for $\pi \in \Pi(n)$ is defined to be the integer partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$ where the number of parts of size i is equal to the number of blocks of size i of π . However, to state the results in a uniform way, we will use the following different definition of $\text{type}(\pi)$.

Definition 2. The *type* of a partition $\pi \in \Pi(n)$, denoted by $\text{type}(\pi)$, is the sequence $(b; b_1, b_2, \dots, b_n)$ where b_i is the number of blocks of π of size i and $b = b_1 + b_2 + \dots + b_n$. The *type* of $\pi \in \Pi_B(n)$ (or $\pi \in \Pi_D(n)$), denoted by $\text{type}(\pi)$, is the sequence $(b; b_1, b_2, \dots, b_n)$ where b_i is the number of unordered pairs $(B, -B)$ of nonzero blocks of π of size i and $b = b_1 + b_2 + \dots + b_n$. For a partition π in either $\Pi(kn)$, $\Pi_B(kn)$ or $\Pi_D(kn)$, if the size of each block of π is divisible by k , then we define the k -*type* $\text{type}^{(k)}(\pi)$ of π to be $(b; b_k, b_{2k}, \dots, b_{kn})$ where $\text{type}(\pi) = (b; b_1, b_2, \dots, b_{kn})$.

Kreweras [10, Theorem 4] proved that the number of $\pi \in \text{NC}(n)$ with $\text{type}(\pi) = (b; b_1, b_2, \dots, b_n)$ is equal to

$$(5) \quad \frac{n!}{b_1!b_2! \dots b_n!(n-b+1)!} = \frac{1}{b} \binom{b}{b_1, b_2, \dots, b_n} \binom{n}{b-1}.$$

Athanasiadis [2, Theorem 2.3] proved that the number of $\pi \in \text{NC}_B(n)$ with $\text{type}(\pi) = (b; b_1, b_2, \dots, b_n)$ is equal to

$$(6) \quad \frac{n!}{b_1!b_2! \dots b_n!(n-b)!} = \binom{b}{b_1, b_2, \dots, b_n} \binom{n}{b-1}.$$

Athanasiadis and Reiner [3, Theorem 1.3] proved that the number of $\pi \in \text{NC}_D(n)$ with $\text{type}(\pi) = (b; b_1, b_2, \dots, b_n)$ is equal to

$$(7) \quad \frac{(n-1)!}{b_1!b_2! \dots b_n!(n-1-b)!} = \binom{b}{b_1, b_2, \dots, b_n} \binom{n-1}{b-1},$$

if $b_1 + 2b_2 + \dots + nb_n \leq n-2$, and

$$(8) \quad (2(n-b) + b_1) \frac{(n-1)!}{b_1!b_2! \dots b_n!(n-b)!} \\ = 2 \binom{b}{b_1, b_2, \dots, b_n} \binom{n-1}{b} + \binom{b-1}{b_1-1, b_2, \dots, b_n} \binom{n-1}{b-1},$$

if $b_1 + 2b_2 + \dots + nb_n = n$.

Armstrong [1, Theorem 4.4.4 and Theorem 4.5.11] generalized (5) and (6) as follows: the number of multichains $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}^{(k)}(n)$ and in $\text{NC}_B^{(k)}(n)$ with $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ are equal to, respectively,

$$(9) \quad \frac{(\ell kn)!}{b_1!b_2! \dots b_n!(\ell kn - b + 1)!} = \frac{1}{b} \binom{b}{b_1, b_2, \dots, b_n} \binom{\ell kn}{b-1},$$

and

$$(10) \quad \frac{(\ell kn)!}{b_1!b_2! \dots b_n!(\ell kn - b)!} = \binom{b}{b_1, b_2, \dots, b_n} \binom{\ell kn}{b}.$$

Krattenthaler and Müller [8] generalized all the above known results except (4) in the following three theorems.

Theorem 1. [8, Corollary 12] *Let $b, b_1, b_2, \dots, b_n$ and $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers satisfying $\sum_{i=1}^n b_i = b$, $\sum_{i=1}^n i \cdot b_i = n$, $\sum_{i=1}^{\ell+1} s_i = n-1$ and $s_1 = n-b$. Then the number of multichains $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ and $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ is equal to*

$$\frac{1}{b} \binom{b}{b_1, b_2, \dots, b_n} \binom{kn}{s_2} \dots \binom{kn}{s_{\ell+1}}.$$

Theorem 2. [8, Corollary 14] *Let $b, b_1, b_2, \dots, b_n$ and $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers satisfying $\sum_{i=1}^n b_i = b$, $\sum_{i=1}^n i \cdot b_i \leq n$, $\sum_{i=1}^{\ell+1} s_i = n$ and $s_1 = n-b$. Then the number of multichains $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}_B^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ and $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ is equal to*

$$\binom{b}{b_1, b_2, \dots, b_n} \binom{kn}{s_2} \dots \binom{kn}{s_{\ell+1}}.$$

Theorem 3. [8, Corollary 16] *Let $b, b_1, b_2, \dots, b_n$ and $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers satisfying $\sum_{i=1}^n b_i = b$, $\sum_{i=1}^n i \cdot b_i \leq n$, $\sum_{i=1}^n i \cdot b_i \neq n-1$, $\sum_{i=1}^{\ell+1} s_i = n$ and $s_1 = n-b$. Then the number of multichains $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}_D^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ and $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ is equal to*

$$\binom{b}{b_1, b_2, \dots, b_n} \binom{k(n-1)}{s_2} \dots \binom{k(n-1)}{s_{\ell+1}},$$



if $b_1 + 2b_2 + \dots + nb_n \leq n-2$, and

$$\begin{aligned} & 2 \binom{b}{b_1, b_2, \dots, b_n} \binom{k(n-1)}{s_2} \dots \binom{k(n-1)}{s_{\ell+1}} \\ & + \frac{s_i - 1}{b - 1} \binom{b-1}{b_1 - 1, b_2, \dots, b_n} \sum_{i=2}^{\ell+1} \binom{k(n-1)}{s_2} \dots \binom{k(n-1)}{s_i - 1} \dots \binom{k(n-1)}{s_{\ell+1}}, \end{aligned}$$

if $b_1 + 2b_2 + \dots + nb_n = n$.

Krattenthaler and Müller's proofs of Theorems 1, 2 and 3 were not combinatorial. Especially, in the introduction, they wrote that Theorems 1 and 2 seem amenable to combinatorial proofs, however, to find a combinatorial proof of Theorem 3 seems rather hopeless. In this paper, we will give combinatorial proofs of Theorems 1, 2 and 3.

This paper is organized as follows. In Section 2 we recall the definition of $\text{NC}_B(n)$ and $\text{NC}_D(n)$. In Section 3 we recall the bijection ψ in [7] between $\text{NC}_B(n)$ and the set of pairs (σ, x) , where $\sigma \in \text{NC}(n)$ and x is either $\emptyset$, an edge or a block of σ . Then we find a necessary and sufficient condition for the two pairs (σ_1, x_1) and (σ_2, x_2) to be $\psi^{-1}(\sigma_1, x_1) \leq \psi^{-1}(\sigma_2, x_2)$ in the poset $\text{NC}_B(n)$. This property will play a crucial role to prove Theorem 3. In Section 4 we prove Theorem 1 by modifying the argument of Edelman [6]. For $0 < r < k$, we consider the subposet $\text{NC}^{(k)}(n; r)$ of $\text{NC}(nk+r)$ consisting of the partitions π such that all but one blocks of π have sizes divisible by k . Then we prove similar chain enumeration results for $\text{NC}^{(k)}(n; r)$. We also prove that the poset $\widetilde{\text{NC}}^{(2k)}(2n+1)$ suggested by Armstrong is isomorphic to $\text{NC}^{(2k)}(n; k)$. With this, we prove Armstrong's conjecture on the zeta polynomial of $\widetilde{\text{NC}}^{(2k)}(2n+1)$ and answer the question on rank-, type-selection formulas [1, Conjecture 4.5.14 and Open Problem 4.5.15]. In Section 5 we prove a

FIGURE 1. The circular representation of $\{\{1, 2, 5, 10\}, \{3, 4\}, \{6, 7, 9\}, \{8\}\}$.FIGURE 2. The circular representations of $\{\pm\{1, 4, -5\}, \pm\{2, 3\}\}$ and $\{\{1, 4, 5, -1, -4, -5\}, \pm\{2, 3\}\}$.

generalization of (4) and Theorem 2. In Section 6 we prove Theorem 3. All the arguments in this paper are purely combinatorial.

2. NONCROSSING PARTITIONS OF CLASSICAL TYPES

Recall that $\Pi(n)$ denotes the poset of partitions of $[n]$ and $\Pi_B(n)$ (resp. $\Pi_D(n)$) denotes the poset of partitions of type B_n (resp. D_n). For simplicity, we will write a partition of type B_n or D_n in the following way:

$$\{\pm\{1, -3, 6\}, \{2, 4, -2, -4\}, \pm\{5, 8\}, \pm\{7\}\},$$

which means

$$\{\{1, -3, 6\}, \{-1, 3, -6\}, \{2, 4, -2, -4\}, \{5, 8\}, \{-5, -8\}, \{7\}, \{-7\}\}.$$

The *circular representation* of $\pi \in \Pi(n)$ is the drawing obtained as follows. Arrange n vertices around a circle which are labeled with the integers $1, 2, \dots, n$. For each block B of π , draw the convex hull of the vertices whose labels are the integers in B . For example, see Figure 1. It is easy to see that the following definition coincides with the definition of a noncrossing partition in the introduction: π is a *noncrossing partition* if the convex hulls in the circular representation of π do not intersect.

Let $\pi \in \Pi_B(n)$. The *circular representation* of π is the drawing obtained as follows. Arrange $2n$ vertices in a circle which are labeled with the integers $1, 2, \dots, n, -1, -2, \dots, -n$. For each block B of π , draw the convex hull of the vertices whose labels are the integers in B . For example, see Figure 2. Note that the circular representation of $\pi \in \Pi_B(n)$ is invariant, if we do not concern the labels, under the 180° rotation, and the zero block of π , if exists, corresponds to the convex hull containing the center. Then π is a *noncrossing partition of type B_n* if the convex hulls do not intersect.




FIGURE 3. The type D circular representations of $\{\pm\{1, -5, -6\}, \pm\{2, 4, -7\}, \pm\{3\}\}$ and $\{\pm\{1, 6, -1, -6\}, \pm\{2, 4, 5\}, \pm\{3\}\}$.




FIGURE 4. The standard representation of $\{\{1, 2, 5, 10\}, \{3, 4\}, \{6, 7, 9\}, \{8\}, \{11\}, \{12, 14\}, \{13\}\}$.




FIGURE 5. The standard representation of $\{\pm\{1, 2, -8\}, \pm\{3, -7\}, \pm\{4, 5\}, \pm\{6\}\}$.

Let $\pi \in \Pi_D(n)$. The *circular representation* of π is the drawing obtained as follows. Arrange $2n - 2$ vertices labeled with $1, 2, \dots, n - 1, -1, -2, \dots, -(n - 1)$ in a circle and put a vertex labeled with $\pm n$ at the center. For each block B of π , draw the convex hull of the vertices whose labels are in B . Then π is a *noncrossing partition of type D_n* if the convex hulls do not intersect in their interiors. For example, see Figure 3.

Let $\pi \in \Pi(n)$. An *edge* of π is a pair (i, j) of integers with $i < j$ such that $i, j \in B$ for a block B of π and there is no other integer k in B with $i < k < j$. The *standard representation* of π is the drawing obtained as follows. Arrange the integers $1, 2, \dots, n$ in a horizontal line. For each edge (i, j) of π , connect the integers i and j with an arc above the horizontal line. For example, see Figure 4. Then π is a noncrossing partition if and only if the arcs in the standard representation do not intersect.

Let $\pi \in \Pi_B(n)$. The *standard representation* of π is the drawing obtained as follows. Arrange the integers $1, 2, \dots, n, -1, -2, \dots, -n$ in a horizontal line. Then connect the integers i and j with an arc above the horizontal line for each pair (i, j) of integers such that i, j are in the same block B of π and there is no other integer in B between i and j in the horizontal line. For example, see Figure 5. Then π is a noncrossing partition of type B_n if and only if the arcs in the standard representation do not intersect.

Let $\text{NC}_B(n)$ denote the subposet of $\Pi_B(n)$ consisting of the noncrossing partitions of type B_n . Note that for $\pi \in \text{NC}_B(n)$, $\text{rank}(\pi) = n - \text{nz}(\pi)$, where $\text{nz}(\pi)$ denotes the number of unordered pairs $(B, -B)$ of nonzero blocks of π .




FIGURE 6. The partition τ obtained from the partition π in Figure 5 by removing all the negative integers. Then $X = \{\{1, 2\}, \{3\}, \{7\}, \{8\}\}$ is the set of blocks of τ which are not blocks of π .




FIGURE 7. The partition σ obtained from the partition τ in Figure 5 by unioning $\{1, 2\}$, $\{8\}$ and $\{3\}$, $\{7\}$ which are the blocks in $X = \{\{1, 2\}, \{3\}, \{7\}, \{8\}\}$. Since X has even number of blocks, x is the edge $(3, 7)$. Then $\psi(\pi) = (\sigma, x)$ for the partition π in Figure 5.

3. INTERPRETATION OF NONCROSSING PARTITIONS OF TYPE B_n

Let $\mathfrak{B}(n)$ denote the set of pairs (σ, x) , where $\sigma \in \text{NC}(n)$ and x is either $\emptyset$, an edge or a block of σ . Note that since for each $\sigma \in \text{NC}(n)$, we have $n + 1$ choices for x with $(\sigma, x) \in \mathfrak{B}(n)$, one may consider $\mathfrak{B}(n)$ as $\text{NC}(n) \times [n + 1]$.

Let us recall the bijection $\psi : \text{NC}_B(n) \rightarrow \mathfrak{B}(n)$ in [7].

Let $\pi \in \text{NC}_B(n)$. Then let τ be the partition of $[n]$ obtained from π by removing all the negative integers and let X be the set of blocks of τ which are not blocks of π . For example, see Figure 6. Now assume that X has k blocks $A_1, A_2, \dots, A_k$ with $\max(A_1) < \max(A_2) < \dots < \max(A_k)$. Let σ be the partition obtained from τ by unioning A_r and A_{k+1-r} for all $r = 1, 2, \dots, \lfloor (k-1)/2 \rfloor$. Let

$$x = \begin{cases} \emptyset, & \text{if } k = 0; \\ (\min(A_{k/2}), \max(A_{k/2+1})), & \text{if } k \neq 0 \text{ and } k \text{ is even;} \\ A_{(k+1)/2}, & \text{if } k \text{ is odd.} \end{cases}$$

Then we define $\psi(\pi)$ to be the pair (σ, x) . For example, see Figure 7.

Theorem 4. [7] *The map $\psi : \text{NC}_B(n) \rightarrow \mathfrak{B}(n)$ is a bijection. Moreover, for $\pi \in \text{NC}_B(n)$ with $\text{type}(\pi) = (b; b_1, b_2, \dots, b_n)$ and $\psi(\pi) = (\sigma, x)$, we have $\text{type}(\sigma) = \text{type}(\pi)$ if π does not have a zero block; and $\text{type}(\sigma) = (b+1; b_1, \dots, b_i+1, \dots, b_n)$ if π has a zero block of size $2i$.*

Now we will find a necessary and sufficient condition for $(\sigma_1, x_1), (\sigma_2, x_2) \in \mathfrak{B}(n)$ to be $\psi^{-1}(\sigma_1, x_1) \leq \psi^{-1}(\sigma_2, x_2)$ in $\text{NC}_B(n)$.

For a partition π (either in $\Pi(n)$ or in $\Pi_B(n)$), we write $i \stackrel{\pi}{\sim} j$ if i and j are in the same block of π and $i \not\stackrel{\pi}{\sim} j$ otherwise. Note that if $\psi(\pi) = (\sigma, x)$, then we have $i \stackrel{\sigma}{\sim} j$ if and only if $i \stackrel{\pi}{\sim} j$ or $i \stackrel{\pi}{\sim} -j$. The following lemmas are clear from the construction of ψ .

Lemma 5. *Let $\psi(\pi) = (\sigma, x)$. Then we have the following.*

- (1) $x = \emptyset$ if and only if there is no integer i with $i \stackrel{\pi}{\sim} -i$.
- (2) x is an edge if and only if there is no integer i with $i \stackrel{\pi}{\sim} -i$ and there are two positive integers i, j with $i \stackrel{\pi}{\sim} -j$. In this case, there is a unique minimal length edge (a, b) satisfying $a \stackrel{\pi}{\sim} -b$ and we have $x = (a, b)$.

- (3) x is a block if and only if there is an integer i with $i \stackrel{\pi}{\sim} -i$. In this case, x is the block consisting of the positive integers i with $i \stackrel{\pi}{\sim} -i$.

Lemma 6. Let $\psi(\pi) = (\sigma, x)$. For any i, j with $i \stackrel{\sigma}{\sim} j$, we have the following.

- (1) If $x = \emptyset$, then $i \stackrel{\pi}{\sim} j$ and $i \not\stackrel{\pi}{\sim} -j$.
- (2) If x is an edge (a, b) of σ , then $i \not\stackrel{\pi}{\sim} j$ and $i \stackrel{\pi}{\sim} -j$ if $i \leq a < b \leq j$; and $i \stackrel{\pi}{\sim} j$ and $i \not\stackrel{\pi}{\sim} -j$ otherwise.
- (3) If x is a block B of σ , then $i \stackrel{\pi}{\sim} j$ and $i \stackrel{\pi}{\sim} -j$ if $\{i, j\} \subset B$; $i \not\stackrel{\pi}{\sim} j$ and $i \stackrel{\pi}{\sim} -j$ if $\{i, j\} \not\subset B$ and $i < \min B \leq \max B < j$; and $i \stackrel{\pi}{\sim} j$ and $i \not\stackrel{\pi}{\sim} -j$ otherwise.

Proposition 7. Let $\psi(\pi_1) = (\sigma_1, x_1)$ and $\psi(\pi_2) = (\sigma_2, x_2)$. Then $\pi_1 \leq \pi_2$ if and only if $\sigma_1 \leq \sigma_2$ and one of the following holds:

- (1) $x_1 = x_2 = \emptyset$,
- (2) x_2 is an edge (a, b) of σ_2 and x_1 is the unique minimal length edge (i, j) of σ_1 with $i \leq a < b \leq j$ if such an edge exists; and $x_1 = \emptyset$ otherwise.
- (3) x_2 is a block of σ_2 , and x_1 is a block of σ_1 with $x_1 \subset x_2$,
- (4) x_2 is a block of σ_2 and x_1 is an edge (i, j) of σ_1 with $i, j \in x_2$.
- (5) x_2 is a block of σ_2 and x_1 is the minimal length edge (i, j) of σ_1 with $i < \min(x_2) \leq \max(x_2) < j$ if such an edge exists; and $x_1 = \emptyset$ otherwise.

Proof. This is a straightforward verification using the above two lemmas. For instance, let us prove a one direction in the second case.

Assume that $\pi_1 \leq \pi_2$ and x_2 is an edge (a, b) of σ_2 . Clearly, we have $\sigma_1 \leq \sigma_2$. For two positive integers i, j with $i < j$, we claim that $i \stackrel{\pi_1}{\sim} -j$ if and only if $i \stackrel{\sigma_1}{\sim} j$ and $i \leq a < b \leq j$.

Let $i \stackrel{\pi_1}{\sim} -j$. Then we have $i \stackrel{\sigma_1}{\sim} j$. Since $\pi_1 \leq \pi_2$, we have $i \stackrel{\pi_2}{\sim} -j$ and by Lemma 5 we get $i \leq a < b \leq j$.

Conversely, let $i \stackrel{\sigma_1}{\sim} j$ and $i \leq a < b \leq j$. Since $\sigma_1 \leq \sigma_2$, we have $i \stackrel{\sigma_2}{\sim} j$ and by Lemma 6 we get $i \not\stackrel{\pi_2}{\sim} j$ and $i \stackrel{\pi_2}{\sim} -j$. On the other hand $i \stackrel{\sigma_1}{\sim} j$ implies $i \stackrel{\pi_1}{\sim} j$ or $i \stackrel{\pi_1}{\sim} -j$, but since $\pi_1 \leq \pi_2$ we cannot have $i \stackrel{\pi_1}{\sim} j$. Thus we get $i \stackrel{\pi_1}{\sim} -j$.

By the claim and Lemma 5, we are done. The other cases can be shown similarly. $\square$

4. k -DIVISIBLE NONCROSSING PARTITIONS OF TYPE A

Let k be a positive integer. A noncrossing partition $\pi \in \text{NC}(kn)$ is k -divisible if the size of each block is divisible by k . Let $\text{NC}^{(k)}(n)$ denote the subposet of $\text{NC}(kn)$ consisting of k -divisible noncrossing partitions. Then $\text{NC}^{(k)}(n)$ is a graded poset with the rank function $\text{rank}(\pi) = n - \text{bk}(\pi)$, where $\text{bk}(\pi)$ is the number of blocks of π .

To prove (1), Edelman [6] found a bijection between the set of pairs $(\mathbf{c}, a)$ of a multichain $\mathbf{c} : \pi_1 \leq \pi_2 \leq \dots \leq \pi_{\ell+1}$ in $\text{NC}^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ and an integer $a \in [n]$ and the set of $(\ell+1)$ -tuples $(L, R_1, R_2, \dots, R_{\ell})$ with $L \in [n]$, $|L| = n - s_1$, $R_i \in [kn]$, and $|R_i| = s_i$ for $i \in [\ell]$. This bijection has been extended to the noncrossing partitions of type B_n [1, 11] and type D_n [3].

In this section we prove Theorem 1 by modifying the idea of Edelman. Let us first introduce several notations.

4.1. The cyclic parenthesization. Let $P(n)$ denote the set of pairs (L, R) of subsets $L, R \in [n]$ with the same cardinality. Let $(L, R) \in P(n)$. We can identify (L, R) with the *cyclic parenthesization* of (L, R) defined as follows. We place a left parenthesis before the occurrence of i for each $i \in L$ and a right parenthesis after the occurrence of i for each $i \in R$ in the sequence $1, 2, \dots, n$. We consider this sequence in cyclic order.

For $x \in R$, the *size* of x is defined to be the number of integers enclosed by x and its corresponding left parenthesis, which are not enclosed by any other matching pair of parentheses. The *type* of (L, R) , denoted by $\text{type}(L, R)$, is defined to be $(b; b_1, b_2, \dots, b_n)$, where b_i is the number of $x \in R$ whose sizes are equal to i and $b = b_1 + b_2 + \dots + b_n$.

Example 1. Let $(L, R) = (\{2, 3, 9, 11, 15, 16\}, \{1, 4, 5, 8, 9, 12\}) \in P(16)$. Then the cyclic parenthesization is the following:

$$(11) \quad 1) (2 (3 4) 5) 6 7 8) (9) 10 (11 12) 13 14 (15 (16$$

Since we consider (11) in the cyclic order, the right parenthesis of 1 is matched with the left parenthesis of 16 and the right parenthesis of 8 is matched with the left parenthesis of 15. The sizes of 5 and 8 in R are 2 and 4 respectively. We have $\text{type}(L, R) = (6; 1, 4, 0, 1, 0, \dots, 0)$.

Let $\overline{P}(n)$ denote the set of elements $(L, R) \in P(n)$ such that the type $(b; b_1, b_2, \dots, b_n)$ of (L, R) satisfies $\sum_{i=1}^n i b_i < n$. Thus we have $(L, R) \in \overline{P}(n)$ if and only if there is at least one integer in the cyclic parenthesization of (L, R) which is not enclosed by any matching pair of parentheses.

We define a map τ from $\overline{P}(n)$ to the set of pairs (B, π) , where $\pi \in \text{NC}(n)$ and B is a block of π as follows. Let $(L, R) \in \overline{P}(n)$. Find a matching pair of parentheses in the cyclic parenthesization of (L, R) which do not enclose any other parentheses. Remove the integers enclosed by these parentheses, and make a block of π with these integers. Repeat this procedure until there is no parenthesis. Since $(L, R) \in \overline{P}(n)$, we have several remaining integers after removing all the parentheses. These integers also form a block of π and B is defined to be this block.

Example 2. Let (L, R) be the pair in Example 1 represented by (11). Note that $(L, R) \in \overline{P}(16)$. Then $\tau(L, R) = (B, \pi)$, where π consists of the blocks $\{1, 16\}$, $\{2, 5\}$, $\{3, 4\}$, $\{6, 7, 8, 15\}$, $\{9\}$, $\{11, 12\}$ and $\{10, 13, 14\}$, and $B = \{10, 13, 14\}$.

Proposition 8. *The map τ is a bijection between $\overline{P}(n)$ and the set of pairs (B, π) , where $\pi \in \text{NC}(n)$ and B is a block of π . Moreover, if $\tau(L, R) = (B, \pi)$, $\text{type}(\pi) = (b; b_1, b_2, \dots, b_n)$ and $|B| = j$, then $\text{type}(L, R) = (b - 1, b_1, \dots, b_j - 1, \dots, b_n)$.*

Proof. Let $\tau(L, R) = (B, \pi)$. It is clear that $\pi \in \text{NC}(n)$. It is sufficient to find the inverse map of τ .

Let $\pi \in \text{NC}(n)$ and B be a block of π . We will find L and R as follows. First we start with $L = R = \emptyset$ and the cyclic sequence P which is initially $1, 2, \dots, n$. If π has more than one block, then let A be a block of π not equal to B such that the elements of A are consecutive in the cyclic sequence P . Note that we can find such a block because π is a noncrossing partition. We add the first and the last integers of A in the cyclic sequence P to L and R respectively. And then we remove the block A from π and remove the integers in A in the cyclic sequence P . Repeat this procedure until π has only one block B . Then the resulting sets L and R satisfy $\tau(L, R) = (B, \pi)$. Thus τ is a bijection.

The ‘moreover’ statement is obvious from the construction of τ . $\square$

We define $P(n, \ell)$ to be the set of $(\ell + 1)$ -tuples $(L, R_1, R_2, \dots, R_\ell)$ such that $L, R_1, R_2, \dots, R_\ell \subset [n]$ and $|L| = |R_1| + |R_2| + \dots + |R_\ell|$. Similarly, we can consider the labeled cyclic parenthesization of $(L, R_1, R_2, \dots, R_\ell)$ by placing a left parenthesis before i for each $i \in L$ and right parentheses $)_{j_1})_{j_2} \dots)_{j_t}$ labeled with $j_1 < j_2 < \dots < j_t$ after i if $R_{j_1}, R_{j_2}, \dots, R_{j_t}$ are the sets containing i among $R_1, R_2, \dots, R_\ell$. For each element $x \in R_i$, the size of x is defined in the same way as in the case of (L, R) . We define the type of $(L, R_1, R_2, \dots, R_\ell)$ similarly to the type of (L, R) .

Example 3. Let $T = (L, R_1, R_2) = (\{2, 4, 5\}, \{2\}, \{2, 6\}) \in P(7, 2)$. Then the labeled cyclic parenthesization of T is the following:

$$(12) \quad 1 \ (2)_1)_2 \ 3 \ (4 \ (5 \ 6)_2 \ 7$$

Then the size of $2 \in R_1$ is 1, the size of $2 \in R_2$ is 3 and the size of $6 \in R_2$ is 2. Thus the type of T is $(3; 1, 1, 0, \dots, 0)$.

Lemma 9. Let $b, b_1, b_2, \dots, b_n$ and $c_1, c_2, \dots, c_\ell$ be nonnegative integers with $b = b_1 + b_2 + \dots + b_n = c_1 + c_2 + \dots + c_\ell$. Then the number of elements $(L, R_1, R_2, \dots, R_\ell)$ in $P(n, \ell)$ with type $(b; b_1, b_2, \dots, b_n)$ and $|R_i| = c_i$ for $i \in [\ell]$ is equal to

$$\binom{b}{b_1, b_2, \dots, b_n} \binom{n}{c_1} \binom{n}{c_2} \dots \binom{n}{c_\ell}.$$

Proof. Let $(L, R_1, R_2, \dots, R_\ell) \in P(n, \ell)$ satisfies the conditions. Let $x_{i1}, x_{i2}, \dots, x_{ia_i}$ be the elements of R_i with $x_{i1} < x_{i2} < \dots < x_{ia_i}$. Let t_{ij} be the size of $x_{ij} \in R_i$.

Then the sequence of t_{ij} ’s is an arrangement of $\overbrace{1, \dots, 1}^{b_1}, \overbrace{2, \dots, 2}^{b_2}, \dots, \overbrace{n, \dots, n}^{b_n}$. Since $(L, R_1, R_2, \dots, R_\ell)$ is completely determined by R_i ’s and t_{ij} ’s, we are done. $\square$

Let $\overline{P}(n, \ell)$ denote the set of $(L, R_1, R_2, \dots, R_\ell) \in P(n, \ell)$ such that the type $(b; b_1, b_2, \dots, b_n)$ of $(L, R_1, R_2, \dots, R_\ell)$ satisfies $\sum_{i=1}^n ib_i < n$.

Using τ , we define a map τ' from $\overline{P}(n, \ell)$ to the set of pairs $(B, \mathbf{c})$, where $\mathbf{c} : \pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ is a multichain in $\text{NC}(n)$ and B is a block of π_1 as follows. Let $P = (L, R_1, R_2, \dots, R_\ell) \in \overline{P}(n, \ell)$. Applying the same argument as in the case of τ to the labeled cyclic parenthesization of P , we get (B_1, π_1) . Then remove all the right parentheses in R_1 from the cyclic parenthesization and their corresponding left parentheses. By repeating this procedure, we get (B_i, π_i) for $i = 2, 3, \dots, \ell$. Then we obtain a multichain $\mathbf{c} : \pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}(n)$. We define $\tau'(P) = (B_1, \mathbf{c})$.

Proposition 10. The map τ' is a bijection between $\overline{P}(n, \ell)$ and the set of pairs $(B, \mathbf{c})$ where $\mathbf{c} : \pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ is a multichain in $\text{NC}(n)$ and B is a block of π_1 . Moreover, if $\tau'(L, R_1, R_2, \dots, R_\ell) = (B, \mathbf{c})$, the rank jump vector of $\mathbf{c}$ is $(s_1, s_2, \dots, s_{\ell+1})$, $\text{type}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ and $|B| = j$, then the type of $(L, R_1, R_2, \dots, R_\ell)$ is $(b - 1; b_1, \dots, b_j - 1, \dots, b_n)$ and $(|R_1|, |R_2|, \dots, |R_\ell|) = (s_2, s_3, \dots, s_{\ell+1})$.

Proof. We will find the inverse map of τ' . This can be done by using a similar argument in [3, 6, 11]. Let $\mathbf{c} : \pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ be a multichain in $\text{NC}(n)$ and B be a block of π_1 . Let B_i be the block of π_i containing B . Now we construct the corresponding labeled cyclic parenthesization as follows.

Let P_ℓ be the labeled cyclic parenthesization $\tau^{-1}(B_\ell, \pi_\ell)$ where all the right parentheses are labeled with ℓ . Then for $i \in [\ell - 1]$ we define P_i to be the labeled

cyclic parenthesization obtained from P_{i+1} by adding the left parentheses, which are not already in P_{i+1} , and the corresponding right parentheses of $\tau^{-1}(B_i, \pi_i)$ and by labeling the newly added right parentheses with i . It is not difficult to check that the map $(B, \mathbf{c}) \mapsto P_1$ is the inverse map of τ' .

The ‘moreover’ statement is obvious from the construction of τ' . $\square$

Theorem 11. *Let $b, b_1, b_2, \dots, b_n$ and $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers satisfying $\sum_{i=1}^n b_i = b$, $\sum_{i=1}^n i \cdot b_i = n$, $\sum_{i=1}^{\ell+1} s_i = n - 1$ and $s_1 = n - b$. Then the number of multichains $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$, $\text{type}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ is equal to*

$$\frac{1}{b} \binom{b}{b_1, b_2, \dots, b_n} \binom{n}{s_2} \cdots \binom{n}{s_{\ell+1}}.$$

Proof. By Lemma 9 and Proposition 10, the number of pairs $(B, \mathbf{c})$, where $\mathbf{c}$ is a multichain satisfying the conditions and B is a block of π_1 , is equal to

$$\sum_{j=1}^n \binom{b-1}{b_1, \dots, b_j-1, \dots, b_n} \binom{n}{s_2} \cdots \binom{n}{s_{\ell+1}} = \binom{b}{b_1, b_2, \dots, b_n} \binom{n}{s_2} \cdots \binom{n}{s_{\ell+1}}.$$

Since there are $b = \text{bk}(\pi_1)$ choices of B for each $\mathbf{c}$, we get the theorem. Note that Lemma 9 states the number of elements in $P(n, \ell)$. However, by the condition on the type, all the elements in consideration are in $\overline{P}(n, \ell)$. $\square$

Now we can prove Theorem 1.

Proof of Theorem 1. Let $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ be a multichain in $\text{NC}^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ and $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$. Then this is a multichain in $\text{NC}(kn)$ with rank jump vector $(kn - 1 - b, s_2, \dots, s_{\ell+1})$ and $\text{type}(\pi_1) = (b; b'_1, b'_2, \dots, b'_{kn})$ where $b'_{ki} = b_i$ for $i \in [n]$ and $b'_j = 0$ if j is not divisible by k . By Theorem 11, the number of such multichains is equal to

$$\frac{1}{b} \binom{b}{b'_1, b'_2, \dots, b'_{kn}} \binom{kn}{s_2} \cdots \binom{kn}{s_{\ell+1}} = \frac{1}{b} \binom{b}{b_1, b_2, \dots, b_n} \binom{kn}{s_2} \cdots \binom{kn}{s_{\ell+1}}.$$

$\square$

4.2. Augmented k -divisible noncrossing partitions of type A . If all the block sizes of a partition π are divisible by k then the size of π must be divisible by k . Thus k -divisible partitions can be defined only on $[kn]$. We extend this definition to partitions of size not divisible by k as follows.

Let k and r be integers with $0 < r < k$. A partition π of $[kn + r]$ is *augmented k -divisible* if the sizes of all but one of the blocks are divisible by k .

Let $\text{NC}^{(k)}(n; r)$ denote the subposet of $\text{NC}(kn + r)$ consisting of the augmented k -divisible noncrossing partitions. Then $\text{NC}^{(k)}(n; r)$ is a graded poset with the rank function $\text{rank}(\pi) = n - \text{bk}'(\pi)$, where $\text{bk}'(\pi)$ is the number of blocks of π whose sizes are divisible by k . We define $\text{type}^{(k)}(\pi)$ to be $(b; b_1, b_2, \dots, b_n)$ where b_i is the number of blocks B of size ki and $b = b_1 + b_2 + \dots + b_n$.

Theorem 12. *Let $0 < r < k$. Let $b, b_1, b_2, \dots, b_n$ and $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers satisfying $\sum_{i=1}^n b_i = b$, $\sum_{i=1}^n i \cdot b_i \leq n$, $\sum_{i=1}^{\ell+1} s_i = n$ and $s_1 = n - b$.*

Then the number of multichains $\mathbf{c} : \pi_1 \leq \pi_2 \leq \cdots \leq \pi_\ell$ in $\text{NC}^{(k)}(n; r)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ and $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ is equal to

$$\binom{b}{b_1, b_2, \dots, b_n} \binom{kn+r}{s_2} \cdots \binom{kn+r}{s_{\ell+1}}.$$

Proof. Let $m = kn + r - \sum_{i=1}^n ki \cdot b_i$. Observe that, as an element of $\text{NC}(kn+r)$, π_1 has type $(b+1; b'_1, b'_2, \dots, b'_{kn+r})$, where $b'_{ki} = b_i$, $b'_m = 1$ and $b'_j = 0$ if $j \neq m$ and $j \neq ki$ for any $i \in [n]$. For given multichain $\mathbf{c}$ satisfying the conditions, let B be the unique block of π_1 whose size is m . By Lemma 9 and Proposition 10, the number of pairs $(B, \mathbf{c})$ is equal to

$$\binom{b}{b'_1, \dots, b'_m-1, \dots, b'_{kn}} \binom{kn+r}{s_2} \cdots \binom{kn+r}{s_{\ell+1}},$$

which is equal to the desired formula. $\square$

Adding up for all the possible nonnegative integers $b_1, b_2, \dots, b_n$ in Theorem 12, we get the following corollary.

Corollary 13. *Let $0 < r < k$. Let $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers with $s_1 + \cdots + s_{\ell+1} = n$. Then the number of multichains in $\text{NC}^{(k)}(n; r)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ is equal to*

$$\binom{n}{s_1} \binom{kn+r}{s_2} \cdots \binom{kn+r}{s_{\ell+1}}.$$

The *zeta polynomial* $Z(P, \ell)$ of a poset P is the number of multichains $X_1 \leq X_2 \leq \cdots \leq X_\ell$ in P . By summing the formula in Corollary 13 for the all $(\ell+1)$ -tuples $(s_1, \dots, s_{\ell+1})$ with $s_1 + \cdots + s_{\ell+1} = n$, we get the zeta polynomial of $\text{NC}^{(k)}(n; r)$.

Corollary 14. *Let $0 < r < k$. The zeta polynomial of $\text{NC}^{(k)}(n; r)$ is the following:*

$$Z(\text{NC}^{(k)}(n; r), \ell) = \binom{n + \ell(kn+r)}{n}.$$

5. k -DIVISIBLE NONCROSSING PARTITIONS OF TYPE B

Let $\pi \in \text{NC}_B(kn)$. We say that π is a *k -divisible noncrossing partition of type B_n* if the size of each block of π is divisible by k .

Let $\text{NC}_B^{(k)}(n)$ denote the subposet of $\text{NC}_B(kn)$ consisting of k -divisible noncrossing partitions of type B_n . Then $\text{NC}_B^{(k)}(n)$ is a graded poset with the rank function $\text{rank}(\pi) = n - \text{nz}(\pi)$, where $\text{nz}(\pi)$ denotes the number of unordered pairs $(B, -B)$ of nonzero blocks of π .

We can prove Theorem 2 using a similar method in the proof of Theorem 1. Instead of doing this, we will prove the following lemma which implies Theorem 2. Note that the following lemma is also a generalization of (4).

For a multichain $\mathbf{c} : \pi_1 \leq \pi_2 \leq \cdots \leq \pi_\ell$ in $\text{NC}_B^{(k)}(n)$, the *index* $\text{ind}(\mathbf{c})$ of $\mathbf{c}$ is the smallest integer i such that π_i has a zero block. If there is no such integer i , then $\text{ind}(\mathbf{c}) = \ell + 1$.

Lemma 15. *Let $b, b_1, b_2, \dots, b_n$ and $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers satisfying $\sum_{i=1}^n b_i = b$, $\sum_{i=1}^n i \cdot b_i \leq n$, $\sum_{i=1}^{\ell+1} s_i = n$ and $s_1 = n - b$. Then the*

number of multichains $\mathbf{c} : \pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}_B^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$, $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ and $\text{ind}(\mathbf{c}) = i$ is equal to

$$\binom{b}{b_1, b_2, \dots, b_n} \binom{kn}{s_2} \dots \binom{kn}{s_{\ell+1}},$$

if $i = 1$, and

$$\frac{s_i}{b} \binom{b}{b_1, b_2, \dots, b_n} \binom{kn}{s_2} \dots \binom{kn}{s_{\ell+1}},$$

if $i \geq 2$.

Proof. Recall the bijection ψ in Section 3. Let $\psi(\pi_j) = (\sigma_j, x_j)$ for $j \in [\ell]$. By Theorem 7, we have $\sigma_j \in \text{NC}^{(k)}(n)$ and $\sigma_1 \leq \sigma_2 \leq \dots \leq \sigma_\ell$. Since $\text{ind}(\mathbf{c}) = i$, x_i is a block of σ_i and x_{i-1} is not a block of σ_{i-1} . By Theorem 7, $x_1, x_2, \dots, x_{i-2}$ and $x_{i+1}, \dots, x_\ell$ are completely determined by respectively x_{i-1} and x_i together with $\sigma_1, \sigma_2, \dots, \sigma_\ell$. Moreover, x_{i-1} is one of the following:

- an edge (u, v) of σ_{i-1} such that $u, v \in x_i$,
- the minimal length edge (u, v) of π_{i-1} with $u < \min(x_i) \leq \max(x_i) < v$ if such an edge exists; and $\emptyset$ otherwise.

Again by Theorem 7, for any choice of $\mathbf{c}' : \sigma_1 \leq \sigma_2 \leq \dots \leq \sigma_\ell$, x_i and x_{i-1} with the above conditions, we can construct the corresponding multichain $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$.

Assume $i = 1$. Then π_1 has a zero block. Let $2t \cdot k$ be the size of the zero block of π_1 . Then $\text{type}(\sigma_1) = (b+1; b_1, \dots, b_t+1, \dots, b_n)$ and the rank jump vector of $\mathbf{c}'$ is $(s_1 - 1, s_2, \dots, s_{\ell+1})$. Thus the number of such multichains $\mathbf{c}'$ is equal to

$$\frac{1}{b+1} \binom{b+1}{b_1, \dots, b_t+1, \dots, b_n} \binom{kn}{s_2} \dots \binom{kn}{s_{\ell+1}},$$

and the number of choices of x_1 is $b_t + 1$. By multiplying these two numbers, we get the formula for the case $i = 1$.

Now assume $i \geq 2$. Then $\text{type}(\sigma_1) = (b; b_1, b_2, \dots, b_n)$ and the rank jump vector of $\mathbf{c}'$ is $(s_1, \dots, s_i - 1, \dots, s_{\ell+1})$. Thus the number of such multichains $\mathbf{c}'$ is equal to

$$(13) \quad \frac{1}{b} \binom{b}{b_1, b_2, \dots, b_n} \binom{kn}{s_2} \dots \binom{kn}{s_i - 1} \dots \binom{kn}{s_{\ell+1}}.$$

Let us fix such a multichain $\mathbf{c}'$. Note that x_i can be any block of σ_i . If x_i is a block B of σ_i , then there are $1 + e(B)$ choices of x_{i-1} , where $e(B)$ is the number of edges (u, v) in σ_{i-1} with $u, v \in B$. Thus there are $\sum_{B \in \sigma_i} (1 + e(B))$ choices for x_i and x_{i-1} . Note that B is a union of several blocks of σ_{i-1} . Let $t(B)$ be the number of blocks of σ_{i-1} whose union is B . Then $e(B) = |B| - t(B)$. Thus the number of choices of x_i and x_{i-1} is equal to

$$\begin{aligned} \sum_{B \in \sigma_i} (1 + e(B)) &= \sum_{B \in \sigma_i} (1 + |B| - t(B)) \\ &= kn - \sum_{B \in \sigma_i} (t(B) - 1) \\ &= kn - (s_i - 1). \end{aligned}$$

The product of this number and (13) is equal to the formula for the case $i \geq 2$. $\square$

5.1. Armstrong's conjecture. Let $\widetilde{\text{NC}}^{(k)}(n)$ denote the subposet of $\text{NC}^{(k)}(n)$ whose elements are fixed under the 180° rotation in the circular representation.

Armstrong [1, Conjecture 4.5.14] conjectured the following. Let n and k be integers such that n is even and k is arbitrary, or n is odd and k is even. Then

$$Z(\widetilde{\text{NC}}^{(k)}(n), \ell) = \binom{\lfloor (k\ell + 1)n/2 \rfloor}{\lfloor n/2 \rfloor}.$$

If n is even then $\widetilde{\text{NC}}^{(k)}(n)$ is isomorphic to $\text{NC}_B^{(k)}(n/2)$, whose zeta polynomial is already known. If both n and k are odd, then $\widetilde{\text{NC}}^{(k)}(n)$ is empty. Thus the conjecture is only for n and k such that n is odd and k is even. In this case, the formula is equivalent to the following:

$$Z(\widetilde{\text{NC}}^{(2k)}(2n+1), \ell) = \binom{n + \ell(2kn + k)}{n}.$$

Theorem 16. *Let n and k be positive integers. Then*

$$\widetilde{\text{NC}}^{(2k)}(2n+1) \cong \text{NC}^{(2k)}(n; k).$$

Proof. Let $\pi \in \widetilde{\text{NC}}^{(2k)}(2n+1)$. Since π is invariant under the 180° rotation, we can consider π as an element of $\text{NC}_B(2kn+k)$. Note that the sum of the block sizes of π is divisible by $2k$ but not by $4k$. Since all the block sizes are divisible by $2k$ and the nonzero blocks appear in pairs, π must have a zero block B of size divisible by $2k$ but not by $4k$.

Now consider the map $\psi : \text{NC}_B(n) \rightarrow \mathfrak{B}(n)$ defined in Section 3. Let $\psi(\pi) = (\sigma, x)$. Since π has the zero block B , by Theorem 4, x is a block whose size is divisible by k but not by $2k$. All the other blocks in σ have sizes divisible by $2k$. Thus $\sigma \in \text{NC}^{(2k)}(n; k)$. Since x is the only block of σ whose size is not divisible by $2k$, the map $\pi \mapsto \sigma$ is a bijection between $\widetilde{\text{NC}}^{(2k)}(2n+1)$ and $\text{NC}^{(2k)}(n; k)$. Moreover, if $\pi < \pi'$ and $\pi' \mapsto \sigma'$, then $\sigma < \sigma'$. Thus this map is in fact a poset isomorphism. $\square$

For $\pi \in \widetilde{\text{NC}}^{(2k)}(2n+1)$, let $\text{type}^{(k)}(\pi) = (b; b_1, b_2, \dots, b_n)$, where $2b_i$ is the number of blocks of π of size ki which are not invariant under the 180° rotation and $b = b_1 + b_2 + \dots + b_n$.

By Theorem 16, we get the following theorems immediately from Theorem 12, Corollaries 13 and 14.

Theorem 17. *Let $b, b_1, b_2, \dots, b_n$ and $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers satisfying $\sum_{i=1}^n b_i = b$, $\sum_{i=1}^n i \cdot b_i \leq n$, $\sum_{i=1}^{\ell+1} s_i = n$ and $s_1 = n - b$. Then the number of multichains $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\widetilde{\text{NC}}^{(2k)}(2n+1)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ and $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ is equal to*

$$\binom{b}{b_1, b_2, \dots, b_n} \binom{2kn+k}{s_2} \dots \binom{2kn+k}{s_{\ell+1}}.$$

Corollary 18. *Let $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers with $s_1 + s_2 + \dots + s_{\ell+1} = n$. Then the number of multichains in $\widetilde{\text{NC}}^{(2k)}(2n+1)$ with rank jump vector*

$(s_1, s_2, \dots, s_{\ell+1})$ is equal to

$$\binom{n}{s_1} \binom{2kn+k}{s_2} \cdots \binom{2kn+k}{s_{\ell+1}}.$$

Corollary 19. [1, Conjecture 4.5.14] *The zeta polynomial of $\widetilde{\text{NC}}^{(2k)}(2n+1)$ is the following:*

$$Z(\widetilde{\text{NC}}^{(2k)}(2n+1), \ell) = \binom{n + \ell(2kn+k)}{n}.$$

6. k -DIVISIBLE NONCROSSING PARTITIONS OF TYPE D

Krattenthaler and Müller [8] found a combinatorial realization of k -divisible noncrossing partitions of type D_n . Let us review their definition.

Let π be a partition of $\{1, 2, \dots, kn, -1, -2, \dots, -kn\}$. We represent π using the annulus with the integers $1, 2, \dots, kn-k, -1, -2, \dots, -(kn-k)$ in the outer circle in clockwise order and the integers $kn-k+1, kn-k+2, \dots, kn, -(kn-k+1), -(kn-k+2), \dots, -kn$ in the inner circle in counter clockwise order. We represent each block B of π as follows. Let $B = \{a_1, a_2, \dots, a_u\}$. We can assume that $a_1, a_2, \dots, a_i$ are arranged in the outer circle in clockwise order and $a_{i+1}, a_{i+2}, \dots, a_u$ are arranged in the inner circle in counter clockwise order for some i . Then we draw curves, which lie inside of the annulus, connecting a_j and a_{j+1} for all $j \in [u]$, where $a_{u+1} = a_1$. If we can draw the curves for all the blocks of π such that they do not intersect, then we call π a *noncrossing partition on the $(2k(n-1), 2k)$ -annulus*.

A k -divisible noncrossing partition of type D_n is a noncrossing partition π on the $(2k(n-1), 2k)$ -annulus satisfying the following conditions:

- (1) If $B \in \pi$, then $-B \in \pi$.
- (2) For each block $B \in \pi$ with u elements, if $a_1, a_2, \dots, a_u$ are the elements of B such that $a_1, a_2, \dots, a_i$ are arranged in the outer circle in clockwise order and $a_{i+1}, a_{i+2}, \dots, a_u$ are arranged in the inner circle in counter clockwise order, then $a_{j+1} \equiv a_j + 1 \pmod{k}$ for all $j \in [u]$, where $a_{u+1} = a_1$.
- (3) If there is a zero block B of π , i.e. $B = -B$, then B contains all the integers in the inner circle.
- (4) If a block B contains only the integers on the inner circle, then B is either $\{kn-k+1, kn-k+2, \dots, kn\}$ or $\{-(kn-k+1), -(kn-k+2), -kn\}$.

Let $\text{NC}_D^{(k)}(n)$ denote the set of k -divisible noncrossing partitions of type D_n . See Figure 8 and 9 for some elements in $\text{NC}_D^{(k)}(n)$.

Let $\pi \in \text{NC}_D^{(k)}(n)$. If a nonzero block B of π contains integers both in the outer circle and in the inner circle, then we call B an *annular block* of π . Let $\overline{\text{NC}}_D^{(k)}(n)$ denote the set of $\pi \in \text{NC}_D^{(k)}(n)$ containing at least one annular block.

Consider an $(\ell+1)$ -tuple $(L, R_1, \dots, R_\ell)$ of sets $L, R_1, \dots, R_\ell \subset [n]$ with $|L| \leq |R_1| + \dots + |R_\ell|$. Let $R = R_1 \uplus R_2 \uplus \dots \uplus R_\ell$ be the disjoint union of $R_1, R_2, \dots, R_\ell$. We regard the elements in L and R as left and right parentheses of $1, 2, \dots, n$ in cyclic order. If $|L| < |R|$, then there are several right parentheses, which do not have the corresponding left parentheses. We call such a right parenthesis *unmatched*. Let R' be the set of unmatched right parentheses. Let f be a function from R' to positive integers. Then we can represent the $(\ell+2)$ -tuple $(L, R_1, \dots, R_\ell, f)$ as a *doubly labeled cyclic parenthesization* as we did in Section 4 together with the additional label $f(x)$ for each unmatched right parenthesis x .




FIGURE 8. Two elements in $\text{NC}_D^{(4)}(5)$. In the left diagram, the integers in the inner circle form two blocks $\pm\{17, 18, 19, 20\}$.




FIGURE 9. Two elements in $\text{NC}_D^{(4)}(5)$.

Example 4. Let $L = \{2, 3, 5, 6, 12, 13\}$, $R_1 = \{4, 8, 12\}$, $R_2 = \{4, 9\}$, $R_3 = \{1, 2\}$ and $R_4 = \{4, 7\}$. Then (L, R_1, R_2, R_3, R_4) is represented by the following parenthesization:

$$(14) \quad 1)_3 (2 (3 4)_1)_2)_4 (5 (6 7)_4 8)_1 9)_2 10 11 (12)_1)_3 (13$$

The unmatched right parentheses are $4 \in R_4$, $9 \in R_2$ and $12 \in R_3$. Let f be the function on these unmatched right parentheses with values 3, 5 and 2 respectively. Then (L, R_1, R_2, R_3, f) is represented by the following doubly labeled cyclic parenthesization:

$$(15) \quad 1)_3 (2 (3 4)_1)_2)_4^3 (5 (6 7)_4 8)_1 9)_2^5 10 11 (12)_1)_3^2 (13$$

Let $P_D(n, \ell)$ denote the set of $(\ell+2)$ -tuples $(L, R_1, \dots, R_\ell, f)$ where $L, R_1, \dots, R_\ell \subset [n]$, $|L| \leq |R_1| + \dots + |R_\ell|$ and f is a function from the unmatched right parentheses to positive integers.

Let $(L, R_1, \dots, R_\ell, f) \in P_D(n, \ell)$. Let $R = R_1 \uplus R_2 \uplus \dots \uplus R_\ell$ and let R' be the set of unmatched right parentheses. For each right parenthesis $x \in R$, we define the *size of x* as follows. If $x \notin R'$, then the size of x is defined to be the number of integers enclosed by x and its corresponding left parenthesis, which are not enclosed by any other pair of parentheses. If $x \in R'$, then the size of x is defined to be $f(x)$ plus the number of integers between x and its previous unmatched right parenthesis which are not enclosed by any pair of parentheses.

Example 5. Let $(L, R_1, R_2, R_3, R_4, f)$ be the element in $P_D(13, 4)$ represented by (15). If we draw only the right parentheses labeled with their sizes, then we get the following:

$$(16) \quad 1)_3^2 \ 2 \ 3 \ 4)_1^2)_2^3 \ 5 \ 6 \ 7)_4^2 \ 8)_1^2 \ 9)_2^6 \ 10 \ 11 \ 12)_1^4)_3 \ 13$$

Note that we can recover (15) from (16).

Let $P_D^{(k)}(n, \ell)$ denote the set of $(\ell + 2)$ -tuples $(L, R_1, \dots, R_\ell, f) \in P_D(kn - k, \ell)$ with the following conditions. As before, $R = R_1 \uplus R_2 \uplus \dots \uplus R_\ell$ and R' is the set of unmatched right parentheses.

- (1) If $|L| = |R|$, then there is at least one integer which is not enclosed by any pair of parentheses.
- (2) If $|L| < |R|$, then $\sum_{x \in R'} f(x) = k$.
- (3) The size of each right parenthesis $x \in R$ is divisible by k .

For $(L, R_1, \dots, R_\ell, f) \in P_D^{(k)}(n, \ell)$, let $\text{type}^{(k)}(L, R_1, \dots, R_\ell, f) = (b; b_1, b_2, \dots, b_n)$, where b_i is the number of right parentheses whose sizes are equal to ki for $i \in [n]$, and $b = b_1 + b_2 + \dots + b_n$. Note that if $|L| = |R|$ then $\sum_{i=1}^n i \cdot b_i \leq n - 2$, and if $|L| < |R|$ then $\sum_{i=1}^n i \cdot b_i = n$.

Lemma 20. Let $b, b_1, b_2, \dots, b_n$ and $c_1, c_2, \dots, c_\ell$ be nonnegative integers with $b = b_1 + b_2 + \dots + b_n = c_1 + c_2 + \dots + c_\ell$ and $\sum_{i=1}^n i \cdot b_i \leq n$. Then the number of elements $P = (L, R_1, \dots, R_\ell, f)$ in $P_D^{(k)}(n, \ell)$ with $\text{type}^{(k)}(P) = (b; b_1, b_2, \dots, b_n)$ and $(|R_1|, |R_2|, \dots, |R_\ell|) = (c_1, c_2, \dots, c_\ell)$ is equal to

$$\binom{b}{b_1, b_2, \dots, b_n} \binom{k(n-1)}{c_1} \dots \binom{k(n-1)}{c_\ell}.$$

Proof. This can be done by the same argument in the proof of Lemma 9. $\square$

We define a map τ_D from $P_D^{(k)}(n, \ell)$ as follows. Let $P = (L, R_1, \dots, R_\ell, f) \in P_D^{(k)}(n, \ell)$. Let $R = R_1 \uplus R_2 \uplus \dots \uplus R_\ell$ and R' be the set of unmatched right parentheses. We will first define $\pi \in \text{NC}_D^{(k)}(n)$ as follows. We place a left parenthesis before each occurrence of i and $-i$ for each $i \in L$ and right parentheses with subscripts $j_1, j_2, \dots, j_t$ after each occurrence of i and $-i$ if i is in $R_{j_1}, R_{j_2}, \dots, R_{j_t}$ with $j_1 < j_2 < \dots < j_t$ in the following infinite cyclic sequence:

$$(17) \quad \dots, 1, 2, \dots, k(n-1), -1, -2, \dots, -k(n-1), 1, 2, \dots$$

Find a matching pair of parentheses which do not enclose any other parenthesis in (17). Remove the pair of matching parentheses and the integers enclosed by them and all of their infinite copies. The removed integers form a block of π . We repeat this procedure until there is no matching pair of parentheses. Then we have the following three possibilities. Now consider the $(2k(n-1), 2k)$ -annulus.

Case 1: There is no unmatched right parenthesis. This happens when $|L| = |R|$. By definition of $P_D^{(k)}(n, \ell)$, there are remaining integers. Then these integers together with the integers in the inner circle form a zero block of π . We define $\tau_D(P)$ to be the set $\{\pi\}$ consisting of π alone.

Case 2: There is an unmatched right parenthesis, and there is no remaining integer. Since the size of each right parenthesis is divisible by k , there must be only one unmatched right parenthesis x with $f(x) = k$. In this case we add two blocks $\pm\{kn - k + 1, kn - k + 2, \dots, kn\}$ to π . We define $\tau_D(P)$ to be the set $\{\pi\}$ consisting of π alone.

Case 3: There are unmatched right parentheses and remaining integers. Let Y be the set of remaining integers. Let $R' = \{x_1, x_2, \dots, x_j\}$ with $x_1 < x_2 < \dots < x_j$. Then the right parentheses coming from R' divide the initial infinite cyclic sequence (17) into the $2j$ blocks $\pm X_i$'s, where

$$\begin{aligned} X_1 &= \{-a_j + 1, \dots, -k(n-1), 1, 2, \dots, a_1\}, \\ X_i &= \{a_{i-1} + 1, a_{i-1} + 2, \dots, a_i\} \quad \text{for } i = 2, 3, \dots, j. \end{aligned}$$

For $i \in [j]$, we claim that $X_i \cap Y \neq \emptyset$. If $X_i \cap Y = \emptyset$, then the unmatched right parenthesis x_i has size $f(x_i)$ which is divisible by k . Since $\sum_{x \in R'} f(x) = k$ and $f(x) > 0$, we get $f(x_i) = k$, $j = 1$ and $Y = \emptyset$, which is a contradiction to the assumption of this case. Thus we have $X_i \cap Y \neq \emptyset$.

Let $A_i = X_i \cap Y$ and $a_i = \max(A_i)$ for $i \in [j]$. We define B_i^+ and B_i^- for $i \in [j]$ as follows. We first define B_j^+ and B_j^- . There is a unique positive integer c_j in the inner circle such that $c_j \equiv a_j + 1 \pmod{k}$. Let B_j^+ (resp. B_j^-) be the set obtained from A_j by adding $f(a_j)$ consecutive integers in the inner circle starting from c_j (resp. $-c_j$) in counter clockwise order. We define B_{j-1}^+ (resp. B_{j-1}^-) to be the set obtained from A_{j-1} by adding the $f(a_{j-1})$ consecutive integers next to the integers used for B_j^+ (resp. B_j^-) in the inner circle. Repeat this until we get B_1^+ and B_1^- . Let π^+ (resp. π^-) be the partition obtained from π by adding $\pm B_1^+, \pm B_2^+, \dots, \pm B_j^+$ (resp. $\pm B_1^-, \pm B_2^-, \dots, \pm B_j^-$). We define $\tau_D(P)$ to be the set $\{\pi^+, \pi^-\}$.

Example 6. Let X be the element in $P_D^{(4)}(16)$ represented by the following:

$$(18) \quad (1 \ 2 \ 3 \ 4)_2 \ (5 \ 6 \ (7 \ 8 \ 9 \ 10)_2 \ 11 \ 12)_1 \frac{1}{2} \ (13 \ 14 \ 15 \ 16)_2$$

Then $\tau_D^{(4)}(X) = \{\pi\}$, where π is the partition represented by the left diagram in Figure 8 (resp. Figure 9). Let Y be the element in $P_D^{(4)}(16)$ represented by the following:

$$(19) \quad 1 \ 2 \ (3 \ 4 \ 5 \ 6)_1 \frac{1}{2} \ 7 \ 8)_1^2 \ 9 \ (10 \ 11 \ 12 \ 13)_2 \ 14 \ 15)_2^1 \ (16$$

Then $\tau_D^{(4)}(Y) = \{\pi^+, \pi^-\}$, where π^+ is the partition represented by the left diagram in Figure 9 and π^- is the partition obtained from π^+ by changing i to $-i$ for each i in the inner circle.

Proposition 21. Let $\pi \in \text{NC}_D^{(k)}(n)$. Then there is a unique $(L, R, f) \in P_D^{(k)}(n, 1)$ such that $\pi \in \tau_D(L, R, f)$.

Proof. The uniqueness is obvious from the construction of τ_D . We find (L, R, f) as follows. Find a nonzero block B of π such that the integers of B in the outer circle of the annulus are consecutive in clockwise order. If B does not have an integer in

the inner circle, then add $|i|$ to L and $|i'|$ to R where i and i' are the first and the last integers among the consecutive integers. If B has an integer in the inner circle, then we only add $|i'|$ to R , and define $f(|i'|)$ to be the number of integers of B in the inner circle. Remove the integers in B and $-B$ in π , and repeat this until π has no nonzero blocks. After finishing this procedure, we get L, R and f . It is easy to see that $\pi \in \tau_D(L, R, f)$. $\square$

Example 7. The unique (L, R, f) 's for the partitions represented by the left and right diagrams in Figures 8 and 9 are the following.

$$(20) \quad (1 \ 2 \ 3 \ 4) \ (5 \ 6 \ (7 \ 8 \ 9 \ 10) \ 11 \ 12) \ (13 \ 14 \ 15 \ 16)$$

$$(21) \quad 1 \ 2 \ 3 \ 4) \ 5 \ 6 \ (7 \ 8 \ 9 \ 10) \ 11 \ 12)^4 \ (13 \ 14 \ 15 \ 16$$

$$(22) \quad 1 \ 2)^1 \ (3 \ 4 \ 5 \ 6) \ 7 \ 8)^2 \ 9 \ (10 \ 11 \ 12 \ 13) \ 14 \ 15)^1 \ (16$$

$$(23) \quad 1 \ 2 \ 3 \ 4 \ 5 \ 6)^1 \ 7 \ 8)^2 \ 9 \ (10 \ 11 \ 12 \ 13) \ 14 \ 15)^3 \ (16$$

Let $\overline{P}_D^{(k)}(n, \ell)$ denote the set of elements $(L, R_1, \dots, R_\ell, f)$ in $P_D^{(k)}(n, \ell)$ such that $|L| < |R_1| + \dots + |R_\ell|$. For $P \in \overline{P}_D^{(k)}(n, \ell)$ and $\epsilon \in \{-1, 1\}$, we will define a multichain $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}_D^{(k)}(n)$ as follows.

Let $P = (L, R_1, \dots, R_\ell, f)$. Consider the doubly labeled parenthesization representing P as shown in (16). Let $P_1 = P$. Then for $2 \leq i \leq n$, P_i is obtained from P_{i-1} by removing all the right parentheses in R_{i-1} and their corresponding left parentheses, if exist. If unmatched right parentheses are removed, then their f values are added to the next unmatched right parentheses in cyclic order, if exist. For instance, if P_1 is

$$1)_2^3 \ 2 \ 3)_1^2 \ 4 \ 5)_1^1 \ 6 \ (7 \ 8)_1)_2^1 \ 9 \ 10)_1^2 \ 11 \ (12)_2)_2,$$

then, in P_2 the f values of $3, 5 \in R_1$ are added to the f value of $8 \in R_2$, and the f value of $10 \in R_1$ is added to the f value of $1 \in R_2$. Thus P_2 is

$$1)_2^5 \ 2 \ 3 \ 4 \ 5 \ 6 \ 7 \ 8)_2^4 \ 9 \ 10 \ 11 \ (12)_2)_2.$$

Now consider the sequence of sets $\tau_D(P_1), \tau_D(P_2), \dots, \tau_D(P_\ell)$. Observe that at least one of these sets contains two elements. Let i be the smallest integer such that $\tau_D(P_i)$ has two elements. Then $\tau_D(P_i) = \{\pi^+, \pi^-\}$. Let $\pi_1, \pi_2, \dots, \pi_{i-1}$ be the unique elements in $\tau_D(P_1), \tau_D(P_2), \dots, \tau_D(P_{i-1})$ respectively. Let $\pi_i = \pi^+$ if $\epsilon = 1$; and $\pi_i = \pi^-$ if $\epsilon = -1$. There is a unique sequence $\pi_{i+1}, \pi_{i+2}, \dots, \pi_\ell$ of elements in $\tau_D(P_{i+1}), \tau_D(P_{i+2}), \dots, \tau_D(P_\ell)$ respectively such that $\pi_i \leq \pi_{i+1} \leq \dots \leq \pi_\ell$. We define $\tau'_D(P, \epsilon)$ to be the multichain $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}_D^{(k)}(n)$. Note that we have $\pi_j \in \overline{\text{NC}}_D^{(k)}(n)$ for at least one $j \in [\ell]$.

Remark 1. Our map τ'_D is a generalization of the map of Athanasiadis and Reiner for $\text{NC}_D(n)$ defined in [3, Proposition 4.3].

Example 8. If P is (18) (resp. (19)), then $\tau'_D(P)$ is the multichain consisting of two elements in Figure 8 (resp. Figure 9).

Proposition 22. *The map τ'_D is a bijection between $\overline{P}_D^{(k)}(n, \ell) \times \{-1, 1\}$ and the set of multichains $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}_D^{(k)}(n)$ with $\pi_j \in \overline{\text{NC}}_D^{(k)}(n)$ for at least one $j \in [\ell]$. Moreover, if $P = (L, R_1, \dots, R_\ell, f)$ and $\tau'_D(P, \epsilon)$ is the multichain $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$, then $\text{type}^{(k)}(P) = \text{type}^{(k)}(\pi_1)$ and $(|R_1|, |R_2|, \dots, |R_\ell|) = (s_2, s_3, \dots, s_{\ell+1})$.*

Proof. We will find the inverse map of τ'_D . Let $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ be a multichain in $\text{NC}_D^{(k)}(n)$ with $\pi_j \in \overline{\text{NC}}_D^{(k)}(n)$ for at least one $j \in [\ell]$. By Proposition 21, there is a unique element $(L'_i, R'_i, f_i) \in P_D^{(k)}(n, 1)$ with $\pi_i \in \tau_D(L'_i, R'_i, f_i)$. Let j be the smallest integer such that $\pi_j \in \overline{\text{NC}}_D^{(k)}(n)$. Then $\pi_j \in \tau_D(L'_j, R'_j, f_j) = \{\pi^+, \pi^-\}$. We define $\epsilon = 1$ if $\pi_j = \pi^+$ and $\epsilon = -1$ otherwise.

Now we define $P = (L, R_1, R_2, \dots, R_\ell, f) \in \overline{P}_D^{(k)}(n, \ell)$ as follows. First we set $L = L'_\ell$ and $R_\ell = R'_\ell$. Then define $R_{\ell-1}$ to be the set of right parentheses in $R'_{\ell-1}$ whose corresponding left parentheses, if exist, are not already in L . And then we add these corresponding left parentheses to L . Repeat this procedure until we get $L, R_1, R_2, \dots, R_\ell$. Note that the unmatched right parentheses of $R = R_1 \uplus R_2 \uplus \dots \uplus R_\ell$ are exactly those of R'_1 . Let $f = f_1$.

It is straightforward to check that the map sending $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ to (P, ϵ) is the inverse map of τ'_D . The ‘moreover’ statement is obvious from the construction of τ'_D . $\square$

Lemma 23. *Let $b, b_1, b_2, \dots, b_n$ and $s_1, s_2, \dots, s_{\ell+1}$ be nonnegative integers satisfying $\sum_{i=1}^n b_i = b$, $\sum_{i=1}^n i \cdot b_i = n$, $\sum_{i=1}^{\ell+1} s_i = n$ and $s_1 = n - b$. Then the number of multichains $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ in $\text{NC}_D^{(k)}(n)$ with rank jump vector $(s_1, s_2, \dots, s_{\ell+1})$ and $\text{type}^{(k)}(\pi_1) = (b; b_1, b_2, \dots, b_n)$ such that $\pi_j \in \overline{\text{NC}}_D^{(k)}(n)$ for at least one $j \in [\ell]$ is equal to*

$$2 \binom{b}{b_1, b_2, \dots, b_n} \binom{k(n-1)}{s_2} \dots \binom{k(n-1)}{s_{\ell+1}}.$$

Proof. By Proposition 22, the number of such multichains is equal to 2 times the number of $P = (L, R_1, \dots, R_\ell, f) \in \overline{P}_D^{(k)}(n, \ell)$ with $\text{type}^{(k)}(P) = (b; b_1, b_2, \dots, b_n)$ and $(|R_1|, |R_2|, \dots, |R_\ell|) = (s_2, s_3, \dots, s_{\ell+1})$. Thus we are done by Lemma 20. Note that by the condition $\sum_{i=1}^n i \cdot b_i = n$, if $\text{type}^{(k)}(P) = (b; b_1, b_2, \dots, b_n)$, then $P \in \overline{P}_D^{(k)}(n, \ell)$ is equivalent to $P \in P_D^{(k)}(n, \ell)$. $\square$

Now we can prove Theorem 3.

Proof of Theorem 3. Let $\pi_1 \leq \pi_2 \leq \dots \leq \pi_\ell$ be a multichain satisfying the conditions with $\pi_j \notin \overline{\text{NC}}_D^{(k)}(n)$ for all $j \in [\ell]$. Let π'_j be the element in $\text{NC}_B^{(k)}(n-1)$ obtained from π_j by removing all the integers in the inner circle. Then $\pi'_1 \leq \pi'_2 \leq \dots \leq \pi'_\ell$ is a multichain in $\text{NC}_B^{(k)}(n-1)$. Let i be the index of this chain, i.e. the smallest integer such that π'_i has a zero block. Then $i = 1$ if and only if $b_1 + 2b_2 + \dots + nb_n \leq n-2$. The rank jump vector is $(s_1, \dots, s_i - 1, \dots, s_{\ell+1})$ and the type is $(b; b_1, \dots, b_n)$ if $i = 1$; and $(b-1; b_1 - 1, \dots, b_n)$ if $i > 1$. By Lemma 15, the number of such multichains is equal to

$$\binom{b}{b_1, b_2, \dots, b_n} \binom{k(n-1)}{s_2} \dots \binom{k(n-1)}{s_{\ell+1}},$$

if $i = 1$, and

$$\frac{s_i - 1}{b - 1} \binom{b - 1}{b_1 - 1, b_2, \dots, b_n} \binom{k(n - 1)}{s_2} \cdots \binom{k(n - 1)}{s_i - 1} \cdots \binom{k(n - 1)}{s_{\ell+1}},$$

if $i \geq 2$.

Now let us count the number of multichains $\pi_1 \leq \pi_2 \leq \cdots \leq \pi_\ell$ with $\pi_j \in \overline{\text{NC}}_D^{(k)}(n)$ for at least one $j \in [\ell]$. We have such a multichain only if $b_1 + 2b_2 + \cdots + nb_n = n$. Using the formula in Lemma 23, we finish the proof. $\square$

REFERENCES

- [1] Drew Armstrong, *Generalized noncrossing partitions and combinatorics of Coxeter groups*, Mem. Amer. Math. Soc. **2002** (2009), no. 949.
- [2] Christos A. Athanasiadis, *On noncrossing and nonnesting partitions for classical reflection groups*, Electron. J. Combin. **5** (1998), R42.
- [3] Christos A. Athanasiadis and Victor Reiner, *Noncrossing partitions for the group D_n* , SIAM J. Discrete Math. **18** (2005), no. 2, 397–417.
- [4] David Bessis, *The dual braid monoid*, Annales scientifiques de l’Ecole normale supérieure **36** (2003), 647–683.
- [5] T. Brady and C. Watt, *Non-crossing partition lattices in finite reflection groups*, Trans. Amer. Math. Soc. **360** (2008), 1983–2005.
- [6] Paul H. Edelman, *Chain enumeration and non-crossing partitions*, Discrete Math. **31** (1980), 171–180.
- [7] Jang Soo Kim, *New interpretations for noncrossing partitions of classical types*, <http://arxiv.org/abs/0910.2036>.
- [8] C. Krattenthaler and T.W. Müller, *Decomposition numbers for finite Coxeter groups and generalised non-crossing partitions*, <http://arxiv.org/abs/0704.0199>.
- [9] Christian Krattenthaler, *Non-crossing partitions on an annulus*, in preparation.
- [10] G. Kreveras, *Sur les partitions non croisées d’un cycle*, Discrete Math. **1** (1972), 333–350.
- [11] V. Reiner, *Non-crossing partitions for classical reflection groups*, Discrete Math. **177** (1997), 195–222.

E-mail address: jskim@liafa.jussieu.fr

URL: <http://liafa.jussieu.fr/~jskim>